

University of Groningen

Indispensable benefit of independent investigator-driven research in a changing clinical trial landscape

Oosting, Sjoukje F; Fehrmann, Rudolf S; de Vries, Elisabeth G E

Published in:
ESMO Open

DOI:
[10.1136/esmoopen-2017-000272](https://doi.org/10.1136/esmoopen-2017-000272)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Oosting, S. F., Fehrmann, R. S., & de Vries, E. G. E. (2017). Indispensable benefit of independent investigator-driven research in a changing clinical trial landscape. *ESMO Open*, 2(4), [e000272]. <https://doi.org/10.1136/esmoopen-2017-000272>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



CrossMark

Indispensable benefit of independent investigator-driven research in a changing clinical trial landscape

Sjoukje F Oosting, Rudolf S Fehrmann, Elisabeth G E de Vries

To cite: Oosting SF, Fehrmann RS, de Vries EGE. Indispensable benefit of independent investigator-driven research in a changing clinical trial landscape. *ESMO Open* 2017;**2**:e000272. doi:10.1136/esmoopen-2017-000272

Received 7 September 2017
Accepted 9 September 2017

[Click here to see the linked article](#)

In the Oxford dictionary, 'Independent' is defined as 'Free from outside control; not subject to another's authority'. Independence can be critical to identify important issues for patients. The authors of the paper 'Safeguarding the future of independent, academic clinical cancer research in Europe for the benefit of patients' are to be applauded for their strong plea for independent investigator-driven research. In addition, they use this paper to draw attention to the Clinical Academic Cancer Research Forum (CAREFOR) platform, raised to address funding and regulatory challenges faced by academic researchers and foster collaborative cancer research by developing novel approaches to do clinical trials.¹

Thanks to industry-driven studies, over the last decades, numerous effective drugs for treatment of patients with cancer have been registered.

Mounting costs leading to limited affordability as well as side effects of these drugs has increased interest in precise insight in the clinical benefit of these drugs for patients. Since 2015, two scales have been made available by, respectively, European Society of Medical Oncology (ESMO) and American Society of Clinical Oncology (ASCO), namely, the ESMO-Magnitude of Clinical Benefit Scale (ESMO-MCBS) and the ASCO Framework to grade the clinical benefit of drugs.^{2,3} Applying these scales made it clear that far from all studies executed to find relevant indications for new drugs had a high yield for the patient. According to Del Paggio and colleagues, only 15% of 277 randomised controlled trials evaluating systemic therapy for colorectal, breast, non-small cell lung and pancreatic cancer published between 2011 and 2015 met ESMO-MCBS thresholds for meaningful clinical benefit.⁴ These authors propose that

investigators, funding agencies, regulatory agencies and industry should adopt more stringent thresholds for meaningful benefit in the design of future randomised clinical trials. Independent research performed by academia might especially allow to focus on relevant clinical benefit.

Furthermore, several highly relevant clinical questions will not be addressed by the pharmaceutical industry. This applies to studies including surgery, radiotherapy, systemic treatment regimens with non-patented drugs and de-escalation studies.

Fortunately, there are already multiple beautiful examples of independent academic studies that indicated their relevance and changed treatment paradigms. Recently, 23 positive Southwest Oncology Group (SWOG) trials sponsored by the National Cancer Institute (NCI) from 1965 to 2012 were analysed in which a total of 12 361 patients were enrolled. It was estimated that 3.34 million life-years were gained from these trials through 2015. The return on investment was US\$125 per life-year gained. The authors concluded that NCI's investment in its cancer cooperative group research programme has provided exceptional value and benefit to the American public.⁵

The European Organisation for Research and Treatment of Cancer (EORTC) Brain Tumour Group study 26951 is an example of a practice changing independent European academic study. This phase III trial demonstrated that adjuvant procarbazine, lomustine and vincristine administered after radiotherapy for anaplastic oligodendroglioma improved survival. This became only clear after long-term survival analysis with a median follow-up of 140 months.⁶

Such long-term follow-up is typically not reported for industry-sponsored studies.



► <http://dx.doi.org/10.1136/esmoopen-2017-000187>

Department of Medical Oncology, University Medical Center Groningen, University of Groningen, Groningen, Netherlands

Correspondence to
Professor Elisabeth G E de Vries;
e.g.e.de.vries@umcg.nl

Funding of trials like these by a European fund or by several National Funds together is essential. There are in addition numerous unsolved clinical questions that could be of major interest for insurance companies and governments. Examples are studies investigating treatment de-escalation strategies or reducing maintenance therapy, in order to limit toxicity and costs while preserving efficacy. For example, the standard of care of patients with recurrent or metastatic head and neck squamous cell carcinoma, is six cycles of platinum-based chemotherapy with cetuximab followed by maintenance therapy with cetuximab until disease progression.⁷ The additive value of cetuximab maintenance treatment is unclear, while it leads to side effects and additional costs. In the Netherlands, costs for just the drug alone are €27690 for an average male patient with a length of 1.80 m and 80 kg body weight on maintenance therapy for 29.9 weeks, which was the median duration of maintenance therapy in the landmark study.⁸ A clinical trial comparing six cycles of chemotherapy plus cetuximab, with and without cetuximab maintenance would therefore be highly relevant. Such a study could be done relatively easy and quickly as an independent academic trial when funded for data management aspects. For studies investigating equivalence of shorter duration of immune checkpoint inhibition for approved indications, cost savings could be even more pronounced.

Moreover, investigator-driven studies could pay deserved attention to patients often excluded for clinical trials even though they could tremendously benefit from a clinical trial such as patients with second or third malignancies and patients with rare tumours or with brain metastases. Happily, there is activity in this field as exemplified by the EORTC 1206 study (ClinicalTrials.gov NCT01969578) in which patients with metastatic androgen receptor positive salivary gland tumours are randomised between hormonal therapy and chemotherapy.

The Systemic Therapy in Advancing or Metastatic Prostate cancer: Evaluation of Drug Efficacy (STAMPEDE) trial is a very elegant and successful example of an ongoing independent academic study in which different pharmaceutical companies participate. Several experimental treatment arms are compared with an ongoing standard of care arm. Ineffective arms are closed early and new treatment arms can be added. The set up and basic structure of this innovative multiarm, multistage platform is funded by Cancer Research UK, Prostate Cancer UK and the Medical Research Council, with contributions of participating pharmaceutical companies. All information is freely accessible on the study website <http://www.stampedetrial.org> which supports participating centres and patients. Already more than 9000 patients have been included and two practice changing papers have been published.^{9 10} An European funding structure for independent academic trials, preferable connected with a cancer research organisation such as EORTC with expertise in conducting

multinational studies, could advance innovative trials after the example of STAMPEDE at the European level.

Not only a new structure for funding of independent clinical cancer research is important, also harmonisation and simplification of regulations will be important. Studies with drugs may become easier in Europe when the EU clinical trial directive is implemented.

In addition, we will have to nurture and educate the next generation of clinical researchers, for example, with workshops such as the *Methods in Clinical Cancer Research (MCCR) Workshop* jointly organised by EOCO, EORTC, American Association for Cancer Research (AACR) and ESMO. Researchers should have basic knowledge concerning data infrastructures created for assembly of so called 'big data', which can reiteratively be used multiple times to answer different research questions leading in potential to relative cheap scientific breakthroughs. It is clear that also patients need insight in this approach as they will be critical partners in bringing new approaches forward.

In the past, in trials only structured clinical data was collected enabling the researchers to assess efficacy of a therapeutic intervention or procedure. Nowadays, however, increasingly additional data such as biospecimen or omics data, and even patient's personal metrics data assembled with wearable technology, are collected to identify, for instance, potential predictive biomarkers for the therapy under investigation. This approach is rapidly expanding, and interest is also focusing on, for example, the ability of omics to predict short-term and long-term side effects. However, this data might dramatically increase in power when studies can be combined and used to answer other relevant questions not directly related to the main research question of the study. A clear example of an answer not related to the main research question is the development of the Response Evaluation Criteria in Solid Tumours (RECIST) for which the imaging and outcome data of patients were merged from industry as well as cooperative group trials. These criteria were developed because changes in tumour burden are frequently used as surrogates of survival/quality of life. In 2000, the RECIST Working Group simplified the 1981 World Health Response Criteria (WHO) 2 after validation in a large data warehouse. In 2009, RECIST was refined (RECIST 1.1) using an even larger warehouse and recently, based on data of 23 259 patients, RECIST criteria for targeted agents were presented.¹¹

Happily, there are now several big initiatives on sharing data. With respect to reuse and merging of data, an important issue is the ownership of data collected in clinical trials. Four potential owners can be identified: the public, the participating patients, the investigators or the study sponsor. In the ideal setting, any anonymous data obtained within clinical trials including industry-sponsored trials should be available to other interested parties in accordance with the findability, accessibility, interoperability, and (re-)usability (FAIR)



principle, while observing ethical, legal and societal constraints.¹² The International Committee of Medical Journal Editors (ICMJE) recently proposed that authors include a plan for data sharing as part of clinical trial registration. Previously, it has been recognised that crediting data generators is a key incentive for data sharing.¹³ A proposal has been put forward to acknowledge persons who initially gathered the data with the concept called data authorship.¹⁴ It states that in order to be cited as a data author, a person must have made substantial contributions to the original acquisition, quality control and curation of the data, be accountable for all aspects of the accuracy and integrity of the data provided and ensure that the available data set follows the FAIR principle. This means that for merging data clinical studies we have to get used to very long author lists and create solutions as done by the Early Breast Cancer Trialists' Collaborative Group.¹⁵ Their studies already showed that pooling of data from multiple trials for so called big-data analyses that can lead to a more accurate and complete view of the complexity of the data and the problem under investigation. Ultimately, this can lead to faster advances in our medical knowledge for the benefit of the patient. Independent clinical researchers should be a major advocate for this data sharing approach.

A structure in Europe, in which we united can perform Independent, Academic Clinical Cancer Research as advocated by Negrouk *et al*,¹ supported by the several European states, national charities, insurance companies, patient organisations and even pharmaceutical companies would be of major benefit for the patient with cancer and society.

Competing interests EGEdV has received research support from Amgen, Roche/Genentech, Chugai Pharma, Synthon, Astra Zeneca, Radius Health, CytomX and Nordic Nanovector made available to the institution.

Provenance and peer review Commissioned; internally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© European Society for Medical Oncology (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Negrouk A, Lacombe D, Cardoso F, *et al*. Safeguarding the future of independent, academic clinical cancer research in Europe for the benefit of patients. *ESMO Open* 2017;2:e000187.
2. Cherny NI, Sullivan R, Dafni U, *et al*. A standardised, generic, validated approach to stratify the magnitude of clinical benefit that can be anticipated from anti-cancer therapies: the European Society for Medical Oncology Magnitude of Clinical Benefit Scale (ESMO-MCBS). *Ann Oncol* 2015;26:1547–73.
3. Schnipper LE, Davidson NE, Wollins DS, *et al*. Updating the American Society of Clinical Oncology value framework: revisions and reflections in response to comments received. *J Clin Oncol* 2016;34:2925–34.
4. Del Paggio JC, Azariah B, Sullivan R, *et al*. Do contemporary randomized controlled trials meet ESMO thresholds for meaningful clinical benefit? *Ann Oncol* 2017;28:157–62.
5. Unger JM, LeBlanc M, Blanke CD. The effect of positive SWOG treatment trials on survival of patients with cancer in the US population. *JAMA Oncol* 2017.
6. van den Bent MJ, Brandes AA, Taphoorn MJ, *et al*. Adjuvant procarbazine, lomustine, and vincristine chemotherapy in newly diagnosed anaplastic oligodendroglioma: long-term follow-up of EORTC brain tumor group study 26951. *J Clin Oncol* 2013;31:344–50.
7. Vermorken JB, Mesia R, Rivera F, *et al*. Platinum-based chemotherapy plus cetuximab in head and neck cancer. *N Engl J Med* 2008;359:1116–27.
8. Vermorken JB, Remenar E, Hitt R, *et al*. Platinum-based chemotherapy (CT) plus cetuximab in recurrent or metastatic squamous cell carcinoma of the head and neck cancer (R/M-SCCHN): 5-year follow-up data for the extreme trial. *J Clin Oncol* 2014;32.
9. James ND, Sydes MR, Clarke NW, *et al*. Addition of docetaxel, zoledronic acid, or both to first-line long-term hormone therapy in prostate cancer (STAMPEDE): survival results from an adaptive, multiarm, multistage, platform randomised controlled trial. *Lancet* 2016;387:1163–77.
10. James ND, de Bono JS, Spears MR, *et al*. Abiraterone for prostate cancer not previously treated with hormone therapy. *N Engl J Med* 2017;377:338–51.
11. Eisenhauer EA, Therasse P, Bogaerts J, *et al*. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45:228–47.
12. Wilkinson MD, Dumontier M, Aalbersberg IJ, *et al*. The FAIR guiding principles for scientific data management and stewardship. *Sci Data* 2016;3:160018.
13. Kalager M, Adami HO, Bretthauer M, *et al*. Recognizing data generation. *N Engl J Med* 2016;374:1898.
14. Bierer BE, Crosas M, Pierce HH. Data authorship as an incentive to data sharing. *N Engl J Med* 2017;376:1684–7.
15. CTSU. EBCTCG: Early Breast Cancer Trialists' Collaborative Group. <https://www.ctsu.ox.ac.uk/research/ebctcg>



Indispensable benefit of independent investigator-driven research in a changing clinical trial landscape

Sjoukje F Oosting, Rudolf S Fehrmann and Elisabeth G E de Vries

ESMO Open 2017 2:

doi: 10.1136/esmoopen-2017-000272

Updated information and services can be found at:
<http://esmoopen.bmj.com/content/2/4/e000272>

These include:

References

This article cites 12 articles, 3 of which you can access for free at:
<http://esmoopen.bmj.com/content/2/4/e000272#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (171)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>